

14785 Preston Road, Suite 550 | Dallas, Texas 75254 Phone: 214 732 9359 | Fax: 972 980 7836

Notice of Independent Review Decision

DATE OF REVIEW: 7/10/2012

IRO CASE #

DESCRIPTION OF THE SERVICE OR SERVICES IN DISPUTE:

Lidoderm 5% Patch Lumbar Spine.

A DESCRIPTION OF THE QUALIFICATIONS FOR EACH PHYSICIAN OR OTHER HEALTH CARE PROVIDER WHO REVIEWED THE DECISION

D.O. Board Certified in Anesthesiology and Pain Management.

REVIEW OUTCOME

Upon independent review the reviewer finds that the previous adverse			
determination/adverse determinations should be:			
Upheld (Agree)			
Overturned (Disagree)			
Partially Overturned (Agree in part/Disagree in part)			

INFORMATION PROVIDED TO THE IRO FOR REVIEW

Document Type	Date(s) - Month/Day/Year
Texas Department of Insurance	6/20/2012
Notice of Case Assignment	
Utilization Review Determinations	5/07/2012-5/24/2012
Clinic	5/16/2012
Formal contest	
Alpha (2) receptors and agonists in pain management. (Article – Source: Department of Anesthesia, Beth Israel Deaconess Medical Center, Boston, Massachusetts)	October 2001
M.D. Letter to the Worker's compensation Division	4/27/2012

PATIENT CLINICAL HISTORY [SUMMARY]:

This is a female who was injured initially on xx/xx/xx. Patient subsequently developed CRPS-2 of the lower extremities, myalgia, myositis, and muscle spasm. Patient is presently on Tizanidine 4mg. tid, Bupernex 0.3 mg. IM qid, and Lidoderm 5% prn.

Patient has been on this regimen for five years, unchanged. Patient reports that the regimen helps and provides a little bit of normalcy in her life, and functions much better with medications. Her pain score without medications is 7-9/10; pain score with medications is 4-5/10.

ANALYSIS AND EXPLANATION OF THE DECISION INCLUDE CLINICAL BASIS, FINDINGS AND CONCLUSIONS USED TO SUPPORT THE DECISION.

Lidoderm 5% Patch Lumbar Spine is medically necessary.



14785 Preston Road, Suite 550 | Dallas, Texas 75254 Phone: 214 732 9359 | Fax: 972 980 7836

Per ODG references, patient should have a Neuropathic component which the patient has (CRPS-2). Patient has been on Trazadone, which is used as an antidepressant and for insomnia. Trazadone inhibits serotonin re-uptake, antagonizes Alpha-1 adrenergic and serotonin receptors. Patient also tried Gabapentin and Lyrica with no apparent improvement with her pain. Since patient has been stable on her treatment regimen for 5 years without any escalation of her medications and she is using Lidoderm on PRN basis with some success, it is very reasonable to continue her Lidoderm.

A DESCRIPTION AND THE SOURCE OF THE SCREENING CRITERIA OR OTHER CLINICAL BASIS USED TO MAKE THE DECISION:

	ACOEM- AMERICAN COLLEGE OF OCCUPATIONAL & ENVIRONMENTAL
_	MEDICINE UM KNOWLEDGEBASE
	AHCPR- AGENCY FOR HEALTHCARE RESEARCH & QUALITY GUIDELINES
	DWC- DIVISION OF WORKERS COMPENSATION POLICIES OR GUIDELINES
	EUROPEAN GUIDELINES FOR MANAGEMENT OF CHRONIC LOW BACK PAIN
	INTERQUAL CRITERIA
	MEDICAL JUDGEMENT, CLINICAL EXPERIENCE AND EXPERTISE IN
	ACCORDANCE WITH ACCEPTED MEDICAL STANDARDS
	MERCY CENTER CONSENSUS CONFERENCE GUIDELINES
	MILLIMAN CARE GUIDELINES
\boxtimes	ODG- OFFICIAL DISABILITY GUIDELINES & TREATMENT GUIDELINES
	PRESSLEY REED, THE MEDICAL DISABILITY ADVISOR
	TEXAS GUIDELINES FOR CHIROPRACTIC QUALITY ASSURANCE & PRACTICE
	PARAMETERS
	TEXAS TACADA GUIDELINES
	TMF SCREENING CRITERIA MANUAL
	PEER REVIEWED NATIONALLY ACCEPTED MEDICAL LITERATURE (PROVIDE A
	DESCRIPTION)
	OTHER EVIDENCE BASED, SCIENTIFICALLY VALID, OUTCOME
	FOCUSED GUIDELINES